

## Barium bronchogram following hydrostatic reduction of intussusception: presentation of a rare complication and brief review of literature

Murugesh Sukumar\*, Deepak Nair, Mathew Varghese and PS Binu

\*Correspondence email: msukumar66@gmail.com

doi: 10.1029/WFSA-D-19-00008

### Abstract

Intussusception is one of the most common paediatric emergencies. Frequently occurs in males under the age of two years. Majority of cases are idiopathic. The accepted initial treatment of uncomplicated intussusception is non-surgical enema reduction using pneumatic or hydrostatic techniques (HR) under fluoroscopic or ultrasound guidance. For HR Barium (Ba) or saline can be used. Complications of HR are rare even though failure to reduce and perforation has been reported. Barium aspiration following HR is hitherto unreported. Ba aspiration in adults and children has been reported following various upper GI contrast studies<sup>1</sup>. We report a case of Ba aspiration following HR of intussusception in a 9 month old child under general anaesthesia. Ba as such is inert, but can remain in lungs for longer period of time and can confuse future imaging studies of respiratory tract.

**Key words:** intussusception; hydrostatic reduction; barium aspiration; barium bronchogram.

### INTRODUCTION

Intussusception refers to the invagination of a part of the intestine into itself. In uncomplicated cases the accepted initial treatment of intussusception is non-surgical enema reduction using pneumatic or hydrostatic techniques under fluoroscopic or USG guidance with or without anaesthesia. For HR barium or saline can be used. Even though ultrasound-guided saline reduction is currently favoured<sup>2</sup> in our institution we still use Ba for HR under fluoroscopy. Complications are rare with HR but rarely failure to reduce, or perforation has been reported<sup>3</sup>. Xie Xiaolong, in a recent study retrospectively reported 637 patients with intussusception<sup>4</sup>. A total of 621 episodes of intussusception were collected for final analysis out of which 62 (9.98%) patients suffered failed reduction. We present a hitherto unreported complication in an infant following HR, and review the literature briefly.

### Case presentation

Intussusception is one of the most common paediatric emergencies. We report a rare complication of hydrostatic reduction under general anaesthesia and review the limited available literature. A 9 month otherwise healthy child weighing 10kg presented with classical signs of intussusception. He was posted for hydrostatic reduction under fluoroscopy/Ga. The baby presented with a history of vomiting, incessant crying and failure to feed. Constipated for one day,

pre-rectal examination revealed blood stained gloved finger. Hydrostatic reduction was attempted using Ba after induction of Ga. As per our anaesthetic protocol, we gave IV Ketamine 2mg/kg anaesthesia was maintained holding mask with sevoflurane in oxygen and nitrous oxide mixture. Reduction was achieved within 10 minutes. Throughout the procedure, the child remained hemodynamically stable with SaO<sub>2</sub> 100%. Sevoflurane and N<sub>2</sub>O were stopped and maintained on 100% O<sub>2</sub> while holding the face mask. Suddenly it became increasingly difficult to ventilate with mask and child started to desaturate, simultaneously white coloured thick secretions were noticed while attempting to insert an airway. Immediate oral suction was done. Intubated and kept on 100% Oxygen. Auscultation revealed bilateral crepitations and wheezes. Baby was put on steroids and inhaled bronchodilators. Postoperative chest X-ray showed extensive opacities suggestive of Ba aspiration. Several attempts were made to suck out Ba through the endotracheal tube but little could be retrieved. A feeding tube was passed to empty the stomach. Bronchoscopy or BAL was not performed. He was shifted to Paediatric ICU for further management. Extubated after 48 hrs un-eventfully. Remained stable after extubation. Baby was discharged on 5th day.

### Post aspiration

X-ray showed well-defined radio-opaque opacities

**Murugesh Sukumar MD**

Khoula Hospital Muscat,  
Muscat,  
OMAN

**Deepak Nair MD DNB**

Ernakulam Medical Centre,  
Kochi  
INDIA

**Mathew Varghese MD DA**

Ernakulam Medical Centre,  
Kochi  
INDIA

**P S Binu FRDS (LOND), FRCS(D),  
DCH (LOND)**

Ernakulam Medical Centre,  
Kochi  
INDIA



**Figure 1:** Barium in lungs X-ray

through all zones of the right lung and but less extensive changes on the left side. Also residual Ba in bowel loops was clearly seen. Feeding tube partially filled with Ba also visible. Ba had accumulated mainly in the Rt lung may be due to the supine position.

### Discussion

In our institution over a period of 10 years we have performed approximately 240 cases of HR using Ba. This was the only case of Ba aspiration following HR during this period. Intussusception typically presents with the sudden onset of colicky but worsening abdominal pain, sometimes with vomiting and bloody stools, alternating with relatively pain-free periods. If left untreated, the bowel can perforate.

### Management choices

For stable patients with evidence of intussusception non-operative reduction is recommended (HR). This procedure involves introducing a substance (air or liquid) into the bowel, via the rectum, with a set pressure that reduces the 'telescoped' bowel into its normal position. The standard method of reduction is to place a reservoir of contrast 100cm above the patient to ensure sustained and constant hydrostatic pressure. Ba (50% dilution of 125w/v Ba.) is used as the contrast agent. Successful reduction is indicated by the free flow of contrast into the small intestine. Very few complications are associated with HR with or without GA. Perforation and failure of reduction has been reported<sup>3,4</sup> Ba regurgitation and aspiration following HR has never been reported in literature.

In our case the standard protocol for Ba reduction was followed regarding the height of Ba container. Here Ba enema successfully reduced the intussusception and may be through an incompetent ileocecal valve entered the large intestine from there to the small intestine and filled the stomach leading to regurgitation and aspiration. We did not anticipate this event and were caught unaware. Several cases of Ba aspiration has been described following upper GI contrast studies. Most of them are usually associated with congenital malformations of gut and airway, like tracheo-oesophageal cleft, tracheo-oesophageal fistula.<sup>5</sup> Ba is inert as such but since it went through the stomach chances of chemical broncho pneumonia was there. In modern era Ba aspiration is typically accidental.<sup>7</sup> Aspiration of significant amounts of Ba in infants is rare and there is no consensus in the literature on how to manage such aspiration. The sequelae of Ba aspiration depends largely on various factors including patient's age, pre-existing lung conditions, the strength of the Ba used<sup>6,7</sup>, the volume aspirated, and the presence of gastric contents<sup>10</sup> the subject's posture at the time of aspiration and respiratory clearance mechanisms such as cough can affect the eventual presence of Ba in the tracheobronchial tree and lungs. Few reports are there in infants. Acute respiratory distress, pneumonitis, sepsis, and even death has been reported in adults<sup>6</sup>. Aspiration of Ba along with gastric contents can give a conflicting picture. Lopez-Castilla et al. reports ARDS like picture and the resultant desaturation in a 2-month-old child with gastro-oesophageal reflux after aspirating Ba following a contrast study<sup>1</sup>. In an attempt to clear the Ba, fiberoptic bronchoalveolar lavages (BAL) were tried. A CT scan of the chest taken 4 months later still showed Ba and micro-nodular densities.

BAL is not routinely recommended as suggested by Wani and Yeola<sup>10</sup>, because of the possibility of further spread of Ba down the lungs. Long-term effects of Ba aspiration provide varying information. Small amounts of Ba are well-tolerated. Residual particles that are not expectorated or removed by mucociliary clearance accumulate in alveolar spaces, using high-resolution CT scans Voloudaki et al, inferred that the Ba particles are likely to be phagocytosed by alveolar macrophages and can potentially cause interstitial fibrosis by crossing into the alveolar or peribronchial interstitial tissue<sup>11</sup> Marchiori et al. differs in this matter in their description of Baritosis in which inhaled particulate matter lies in the lungs for years without producing symptoms or significant respiratory impairment<sup>12</sup>. A study in dogs by Wilson et al. only a mild transient inflammatory reaction following Ba bronchography, which was quickly replaced by a harmless foreign body reaction<sup>13</sup>. The same investigators did not find any ill-effects, either acute or chronic. They concluded that Ba sulfate in the lung behaves as a relatively inert foreign body. Various studies by different investigators involving Ba bronchography has not resulted in short or long term ill effects<sup>14</sup>. The innocent nature of Ba in lungs is provided by Doig who described nine cases of Baritosis in factory workers exposed to Ba dust<sup>15</sup>. He described it as a benign pneumoconiosis. He followed up these subjects for 9 years and reported partial clearing of radiographic changes.

Before the advent of computed tomography and bronchoscopy Ba had been used as a contrast medium for bronchography. Although historically found to be safe in this context, aspiration of Ba may also produce a chemical pneumonitis in some cases.

Limited data exist characterizing the long-term prognoses of patients after isolated Ba aspiration events. Most patients appear to have complete recovery given the inert nature of Ba; however, high-resolution chest imaging has detected subtle evidence of early fibrosis even 1 year after aspiration<sup>11</sup>. Wani and Yeola took the contrary view and recommended against bronchoalveolar lavage arguing that it may disseminate the Ba further within the bronchoalveolar system<sup>10</sup>.

In paediatrics, imaging techniques may not differentiate primary from secondary aspiration. Factors that are likely to influence outcomes post-aspiration are the child's pre-existing clinical state, the volume, and concentration of the Ba sulfate used as well as the immediate post-aspiration clinical status. If Ba is aspirated, immediate management of Ba aspiration involves an attempt to suck out Ba as much as possible from the tracheobronchial tree and supportive care. Antibiotics and steroids are not indicated routinely. Ba aspiration is rare and often produces dramatic radiographic findings, but is generally associated with a favourable prognosis. Evidence favours supportive care in most cases, with therapeutic bronchoalveolar lavage to be considered only in cases with significant respiratory symptomatology<sup>5</sup>. The evidence suggests that the Ba will remain in the lung for an extended time, be relatively inert and that the risk of fibrosis or other complications remains low.

### Conclusion

Ba aspiration has been reported following several Ba studies. Since Ba can remain in lungs for an indefinite period of time and can interfere with future radiological studies parents were warned about this. Literature review did not reveal Ba aspiration following hydrostatic reduction of intussusception in a child. In the above case after iv induction, anaesthesia was maintained holding mask. We still continue the same technique. However, endotracheal intubation and protection of airway should be seriously considered we hereby report this rare complication and alert the concerned physicians about this potentially fatal complication.

### REFERENCES

1. Lopez-Castilla J, Cano M, Munoz M, Soult JA, Andrés A, Montilla M, et al. Massive bronchoalveolar aspiration of Ba sulphate during a radiologic study of the upper digestive tract. *Pediatr Pulmonol* 1997; **24(2)**: 126–7.
2. Kim YG, Choi BI, Yeon KM, Kim JW. Diagnosis and treatment of childhood intussusception using real time ultrasonography and saline enema: preliminary report. *J Korean Soc Med Ultrasound*. 1982; **1**: 66-70.
3. Bramson RT, Blickman JG. Perforation during hydrostatic reduction of intussusception: proposed mechanism and review of the literature. *J Pediatr Surg*. 1992; **27(5)**: 589-91
4. Xiaolong X, Yang W, Qi W, Yiyang Z, Bo X. Risk factors for failure of hydrostatic reduction of intussusception in pediatric patients: A retrospective study. *Medicine (Baltimore)*. 2019; **98(1)**: e13826.
5. Jackson M, Kapur N, Goyal V, Choo K, Sarikwal A, Masters IB et al. Barium aspiration in an infant: a case report and review of management. *Front Pediatr*. 2014; **2**: 37.
6. Taman I, Kortsik C. Severe Ba sulphate aspiration in the lung: clinical presentation, prognosis and therapy. *Respiration* 1999 **66**: 81–4.
7. Whiting J. Aspiration of Barium. *N Engl J Med* 2003; **348**: 2582–310
8. Fruchter O, Dragu R. A deadly examination. *N Engl J Med*; 2003; **348(11)**: 1016
9. Tamm I, Kortsik C. Severe barium sulphate aspiration into the lung: clinical presentation, prognosis and therapy. *Respiration* 1999; **66**: 81–4.
10. Wani B, Yeola M. Aspiration of Ba sulphate in swallow study. *Internet J Pulm Med*; 2008: 10(2).
11. Voloudaki A, Ergazakis N, Gourtsoyiannis N. Late changes in Ba sulphate aspiration: HRCT features. *Eur Radiol*; 2003; **13(9)**: 2226-9.
12. Marchiori E, Souza A, Franquet T, Muller N. Diffuse high-attenuation pulmonary abnormalities: a pattern-oriented diagnostic approach on high-resolution CT. *Am J Roentgenol*; 2005; **184**: 273– 8210.
13. Wilson J, Rubin P, McGee T. The effects of Ba sulfate on the lungs: a clinical and experimental study. *Am J Roentgenol Radium The Nucl Med*; 1959; **82**: 84.
14. Shook D, Felson B. Inhalation bronchography. *Chest*; 1970; **58(4)**: 333–710.
15. Doig A. Baritosis: a benign pneumoconiosis. *Thorax*; 1976; **31**: 30 810.

# Update in Anaesthesia

---

## Guide for Contributors

---

### MISSION OF UIA

*Update in Anaesthesia* (UIA) is an official journal of the World Federation of Societies of Anaesthesiologists (WFSA). UIA's primary goal is to provide high-quality, peer-reviewed, clinically-relevant educational articles aimed at improving the quality and safety of anaesthesia care around the world.

UIA welcomes manuscripts focused on, and not limited to, anaesthesia that is relevant to resource-limited settings. UIA also serves to share knowledge from affiliated associations such as the World Health Organization (WHO) and surgical societies and educators that recognise safe anaesthesia as an essential part of safe surgery.

Over 16,000 anaesthetists around the world access UIA on the internet at <http://www.wfsahq.org/resources/update-in-anaesthesia>. UIA is read by many others including anaesthesia trainees, nurses, surgeons and medical students.

Manuscripts should be submitted to:

<https://www.editorialmanager.com/wfsa/default.aspx>

### GENERAL CONSIDERATIONS FOR PREPARATION AND SUBMISSION OF A MANUSCRIPT

A major focus of UIA is to provide educational materials for those working with limited resources. The types of manuscripts that are published in the journal reflect this. If you have an idea for an article and would like to discuss this further, please email the Editor-in-Chief with a proposed title and a brief (200 word) summary of the scope and content.

Manuscripts are screened by the editorial staff and are sent to independent reviewers who are not informed of the identity of the author(s). The reviewers' comments and suggested revisions will be forwarded to the corresponding author. Manuscripts will be reviewed with plagiarism detection software. Manuscripts may not have been published in whole or any part in another journal.

Some readers of UIA may not use English as their first language. It is therefore important to keep text straightforward and to avoid long sentences and complex terminology. Explain words and abbreviations that may not be universally standardised.

Aim to include the full range of therapies available worldwide. Provide detailed descriptions of therapies available in resource-limited settings. When appropriate, discuss older medications, such as Halothane and Thiopentone that may still be used around the world.

Authors must acknowledge any financial support received or conflict of interest related to the contents of a submitted manuscript.

Contributions to the, 1) clinical care or research described in a manuscript or 2) preparation of a manuscript made by persons other than the authors may be acknowledged.

Upon acceptance for publication copyright becomes vested with the journal.

Manuscripts based on clinical investigation of human subjects must adhere to the ethical principles of the Declaration of Helsinki developed by the World Medical Association. <https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/>

Manuscripts based on clinical investigation of human subjects must include the consent of the patients or study participants or appropriate person (parent/guardian) and a statement of approval from an appropriate Institutional Review Board (IRB)/Ethics Committee. The IRB/Ethics Committee must review and provide oversight or grant exemption from requiring patients' or study participants' consent for the performance of audits and this should be referred to in the text.

Manuscripts must protect patients and study participants from being identified. UIA adheres to the USA HIPAA Privacy Rule that defines 18 identifiers that must be removed to assure the protection of personal health information. [https://privacyruleandresearch.nih.gov/pdf/HIPAA\\_Privacy\\_Rule\\_Booklet.pdf](https://privacyruleandresearch.nih.gov/pdf/HIPAA_Privacy_Rule_Booklet.pdf) (Specifically refer to page 10)

Manuscripts should adhere to Uniform Requirements for Manuscripts Submitted to Biomedical Journals prepared by the International Committee of Medical Journal Editors. <http://www.icmje.org/recommendations/browse/manuscript-preparation/preparing-for-submission.html>

### Manuscript Types

As an official journal of the WFSA, UIA is the appropriate forum for publication of articles describing anaesthesia patient care, original research and audit conducted in, and specifically relevant to, low and middle income resource settings.

Manuscripts describing clinical care, research, procedures, techniques or equipment adapted by anaesthetists to the limited resource setting in which they care for patients are welcome in any of the following formats:

- Review article - Basic Science or Clinical (3000 words maximum)
- Original Research (3000 words maximum)
- Case Report/Clinical Overview (1500 words maximum)
- UIA Short Reports (750 words maximum)
- Editorials (1200 words maximum) - Editorials are usually by invitation of the Editor-in-Chief. Proposals for Editorials may be sent to the Editor-in-Chief for consideration and potential invitation.
- Review of Books or other Educational Materials (eg web app) (750 words maximum)



- Letters to the Editor (1000 words maximum) - Correspondence is welcome on any topic, article or editorial that has appeared in UIA or is of interest to individuals providing anaesthesia patient care.

## COMPONENTS OF THE MANUSCRIPT

Title page that includes:

- Title of the article
- Name and affiliation of each author
- Email, postal address and telephone contact details of the corresponding author
- Key words using terms from the Medical Subject Headings (MeSH) of Index Medicus <https://www.nlm.nih.gov/mesh>
- Abstract that includes no more than 200 words

For research manuscripts note Background, Methods, Results, and Conclusions

## ILLUSTRATIONS / FIGURES

Tables, figures, graphs and illustrations should be submitted as electronic versions of drawings (black on white) or as black and white or color picture files in JPEG format. If computer or internet access is unavailable, print versions may be submitted. If facilities to produce drawings are unavailable, contact the Editor-in-Chief for assistance. Text accompanying tables, figures, graphs and illustrations should be supplied on a page separate from the image.

Tables, figures, graphs and illustrations copied from other publications must be accompanied by a statement that permission for reproduction has been obtained from the publisher. A footnote acknowledging permission to reproduce must accompany the copied table, figure, graph or illustration and the full reference of the original source included in the reference list.

If patients appear in an identifiable photo, a copy of their consent to the use of their image must be submitted.

## REFERENCES

A minority of UIA readers have extensive access to journals. References, therefore, should in general be limited to those that are considered 'desirable supplemental reading', and should not exceed 25 references. Number the references in the order they appear in the text, using the reference number as a superscript at the relevant point in the text.

Reference style should adhere to the format set forth in the Uniform Requirements for Manuscripts Submitted to Biomedical Journals prepared by the International Committee of Medical Journal Editors <http://www.icmje.org>

Abbreviations for journal titles should follow Index Medicus format <https://www.ncbi.nlm.nih.gov/nlmcatalog/journals>

Examples of the recommended format for references are noted below and can be found at [https://www.nlm.nih.gov/bsd/uniform\\_requirements.html](https://www.nlm.nih.gov/bsd/uniform_requirements.html)

1. Reynolds F, O'Sullivan G. Lumbar puncture and headache. 'Atraumatic needle' is a better term than 'blunt needle'. *Br Med J* 1998; 316: 1018.
2. Costigan SN, Sprigge JS. Dural puncture: the patients' perspective. A patient survey of cases at a DGH maternity unit 1983-1993. *Acta Anaesthesiol Scand* 1996; 40: 710-14.
3. Spriggs DA, Burn DJ, French J, Carlidge NE, Bates D. Is bedrest useful after diagnostic lumbar puncture? *Postgrad Med J* 1992; 68: 581-3.

References to textbooks should give book title, place of publication, publisher and year; those of multiple authorship should also include chapter title, first and last page numbers, and names and initials of editors. For example:

1. Roberts F. Chapter 22: Ear, nose and throat surgery. In: Allman KG, Wilson IH, eds. *Oxford handbook of Anaesthesia* (1st edition) Oxford: Oxford University Press, 2001: 506-39.

Unpublished manuscripts in preparation or submitted yet not accepted for publication and personal communications and unpublished observations should be referred to as such in the text.

Articles accepted and not yet published should be included in the references, with the abbreviated journal name, followed by '(in press)'.

## MEDICATIONS

Use international units, e.g. mg.kg<sup>-1</sup> rather than mg/kg. Use SI notation for g, mg, mcg etc. Use internationally accepted non-proprietary generic medication names, e.g. furosemide, epinephrine and avoid trade names.

Manuscripts should be submitted to:

<https://www.editorialmanager.com/wfsa/default.aspx>

UIA's Editorial Team will be delighted to help with the preparation of articles. Please contact the Editors-in-Chief for any assistance.

- Christina Lundgren  
Chris.Lundgren@wits.ac.za
- Victoria Howell  
updateinanaesthesia@gmail.com

If you are unable to upload your manuscript to the online submission system, please email them directly to the editors.

Alternatively manuscripts may be sent to:

World Federation of Societies of Anaesthesiologists (WFSA)  
Dean Bradley House,  
52 Horseferry Rd,  
London  
SW1P 2AF  
England